READ SAFETY DIRECTIONS BEFORE OPENING OR USING



A gel bait for the treatment of cockroach infestations

GENERAL INSTRUCTIONS

To apply the gel, remove the cap on the nozzle and touch the tip to the surface to be treated and slowly depress plunger until sufficient gel is dispensed. Recap the dispenser after treatment is completed.

Where maintenance treatments form part of the professional pest controllers contract it is recommended that cockroach activity be monitored where gel has been applied one month after initial treatment or at more regular intervals in areas of high infestation.

Insecticide Resistance Warning

For insecticide resistance management Maxforce Gold Gel is a Group 2C insecticide.

Some naturally occurring insect biotypes resistant to Maxforce Gold Gel and other Group **2C** insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Maxforce Gold Gel or other Group **2C** insecticides are used repeatedly. The effectiveness of Maxforce Gold Gel on resistant individuals could be significantly reduced. Since the occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience accepts no liability for any losses that may result from the failure of this product to control resistant insects.

Maxforce Gold Gel Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier or Bayer CropScience representative.

PROTECTION OF LIVESTOCK, WILDLIFE, AND OTHERS

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT apply this product in any areas easily accessible to children or animals.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated locked place out of the reach of children. Do not store in direct sunlight and do not store at temperatures exceeding 35°C. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in the garbage.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

EXCLUSION OF LIABILITY

This product as supplied is of a high grade and suitable for the purpose for which it is expressly intended and must be used in accordance with the directions. The user must monitor the performance of any product as climatic, geographical or biological variables and/or developed resistance may affect the results obtained. No responsibility is accepted in respect of this product, save for those non-excludable conditions implied by the Trade Practices Act or any State or Federal legislation.

Maxforce[®] is a registered trademark of Bayer

NRA Approval No.: 55553/0802

FOR 24 HOUR SPECIALIST ADVICE IN EMERGENCY ONLY PHONE 1800 033 111



DIRECTIONS FOR USE

DO NOT place in areas that are routinely washed.

SITUATION	PEST	RATE	CRITICAL COMMENTS
Domestic, Commercial	German and other	Apply at least 2 spots per m ² .	Conduct a thorough inspection of the
and Public Service	common	Increasing the number of spots	premises and then place gel in areas
buildings including; farm	cockroaches.	(up to 5/m ²) may be required	near cockroach harbourages or in
buildings, food processing		for heavy infestations or if the	sheltered areas where signs of
establishments, factories,		American cockroach is the	cockroaches are evident.
hospitals, homes,		main pest.	
kitchens, offices,			Replenish gel if previous application
restaurants, retail outlets,		Recommended spot size is	has been consumed and cockroaches
store-rooms, vehicles and		approximately 5 mm diameter	remain a problem; increasing number
other situations which fall		(0.1 g)	of spots if necessary.
within these definitions.			

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.